

**INTERNATIONAL JOURNAL OF ADVANCES IN
PHARMACY, BIOLOGY AND CHEMISTRY**

Research Article

**Larvicidal activity of Methanolic Extract of the leaves
of *Citrullus lanatus***

Ajithadas Aruna*, Vijayalakshmi K, Karthikeyan V.

College of Pharmacy, Madurai Medical College, Madurai-20, Tamil Nadu, India - 625020.

ABSTRACT

Extensive use of synthetic and chemical insecticides to control mosquitoes result in environment hazards and development of resistance in vector species. This research work is considered, herbals as safe to environment and non-target species and also bio-degradable. To find out larvicidal efficacy of Methanolic extract of *Citrullus lanatus* (MECL) leaves against *Anopheles stephensi*, *Culex quinquefasciatus* and *Aedes aegypti*.

LC₅₀ and LC₉₀ of MECL were determined against *Anopheles stephensi*, *Culex quinquefasciatus* and *Aedes aegypti* by using Abbott's formula. LC₅₀ and LC₉₀ of MECL against *Anopheles stephensi* were found to be 84.23ppm & 989.396ppm. LC₅₀ and LC₉₀ of MECL against *Culex quinquefasciatus* were found to be 51.31ppm & 405.88ppm. LC₅₀ and LC₉₀ of MECL against *Aedes aegypti* were found to be 2645ppm & 138326ppm. Methanolic extract of *Citrullus lanatus* showed promising larvicidal activity against important vectors of Malaria, Filariasis, Dengue and Yellow fever.

KEYWORDS: *Citrullus lanatus*, Larvicidal activity, LC₅₀, LC₉₀.

INTRODUCTION

Mosquito borne diseases are one of the most public health issues in the developing countries. Malaria and other vector-borne diseases contribute to the major disease burden in India. One of the methods to control these diseases is to control the vectors for the interruption of disease transmission. In the past, synthetic organic chemical insecticides based intervention measures for the control of insect pests and disease vectors have resulted in development of insecticide resistance in some medically important vectors of malaria, filariasis, Japanese encephalitis, dengue, hemorrhagic fever, chikungunya and yellow fever transmitted by mosquitoes which cause millions of death every year¹. Hence controlling the mosquitoes in larval stage is important and effective for preventing the above infections^{2,3}. During the last decade, various studies on natural plant products against mosquito vectors indicate them as possible alternatives to synthetic chemical insecticides. Mosquito control programmes largely target the larval stage at their breeding sites with larvicides⁴. Larviciding is a successful method of reducing mosquito population in their breeding place before they emerge as adults. The screening of local

medicinal plants for mosquito larvicidal activity may eventually lead to their use in natural product-based mosquito abatement practices⁵. Plants are well known to contain a complex of chemicals with bioactive potential like deterrents or attractants. Plant extracts are reported to be eco-friendly mosquito control agents^{6,7}.

Citrullus lanatus is well known as Watermelon plant (Family - Cucurbitaceae). Water melon is popular in indigenous system of folk medicine. It is a trailing annual plant with several herbaceous, firm and stout stems^{8,9}. The leaves of *Citrullus lanatus* is used as anti-inflammatory, analgesic, gonorrhoea, mosquitocidal and anti microbial property¹⁰.

Leaves are available throughout the year could be easily collected without any additional cost. Therefore, leaves extracts could be used as a larvicidal agent in an integrated vector control program. The plant *Citrullus lanatus* has been selected (specially the leaves) for the present investigation on the basis of the ethnomedical information and the review of literature as the plant is widely cultivated throughout India.

Larvicidal effect of *Citrullus colocynthis* leaves was previously reported¹¹. This present work has attempted to study mosquito larvicidal property of *C.lanatus* against three mosquito species - *Anopheles stephensi*, *Culex quinquefasciatus* and *Aedes aegypti*.

MATERIALS AND METHODS

Collection and preparation of extract: The leaves of *Citrullus lanatus* were collected in Thuvankurichi during the month of August 2013. The plant specimen was identified and authenticated as '*Citrullus lanatus*' (Cucurbitaceae) by Dr. Stephen, Senior Lecturer in Botany and Taxonomist, Dept. of Botany, The American College, Madurai, Tamil Nadu, India. The authenticated herbarium of plant has been kept in the Department of Pharmacognosy, Madurai Medical College, Madurai. The leaves were washed thoroughly and dried in shade. The shade dried leaves were powdered and used for further studies. Extraction of leaves of *Citrullus lanatus* was carried out by washing the plants and drying at room temperature in 14 days. After that, they were filtered with sieve analyzer to get homogeneous particles and defatted with 2.5L of petroleum ether (60-80°C) by cold maceration method for 72hr. The solvent was then removed by filtration and the marc was dried. The dried marc was re-soaked with 2.5L of Methanol. The steps were performed three times and the combined filtrates were evaporated to a cohesive mass using rota vapour.

Collection of mosquito: The mosquito larvae (*Anopheles stephensi*, *Culex quinquefasciatus* and *Aedes aegypti*) were obtained from ICMR, Madurai.

Preparation of stock solution of Methanolic extract of *C.lanatus*

A stock solution of 100mg/ml of the MECL was prepared by dissolving the required quantity of the extract in distilled water.

Larvicidal activity of MECL¹²⁻¹⁶

The larvicidal activity of plant extract was carried out on late 3rd and early 4th instar larvae of *Anopheles stephensi*, a primary vector of urban malaria, *Culex quinquefasciatus*, a common vector of filariasis, *Aedes aegypti*, common vector of dengue and yellow fever. Twenty larvae were released in 500ml beaker containing 200ml distilled water with varying concentrations of plant extract. The larvae were provided with dog biscuit and yeast powder in a ratio of 3:2 as nutrients. The experiments were carried out at room temperature (26°C ± 2°C). Three replicates of each concentration were run under the same microclimate conditions along with untreated control. The mortality was monitored for 24hr.

Data analysis¹⁷⁻¹⁹

LC₅₀ (lethal concentration to cause 50% mortality in the population) and LC₉₀ (lethal concentration to cause 90% mortality in the population) were determined by plotting the regression line. The percentage mortality was calculated using Abbot's formula and the data so obtained was analyzed by probit analysis by using the software Minitab-15 for dose and time mortality regression lines. The purpose of the probit transformation is to straighten the line so we can estimate LC₅₀ more easily.

$$\text{Percentage Mortality} = \frac{\% \text{Mortality in treated} - \% \text{Mortality in control}}{100 - \% \text{Mortality in control}} \times 100$$

RESULTS AND DISCUSSION

Larvicidal activity of MECL against *Anopheles stephensi*

The results obtained for the larvicidal effect of MECL against *Anopheles stephensi* & the graphical representation were presented (Table 1 & Fig. 1).

From the Table 1 it can be observed that a mortality of 100.00 ± 0.00 was observed for *Anopheles stephensi*. The LC₅₀ and LC₉₀ values were calculated using Probit analysis. The percentage mortality calculated using Abbott's formula versus log concentration was plotted and Y=50 is substituted in the resulting linear equation to obtain the X value. The linear regression equation was found to be y = 37.387x - 21.988 for activity against *Anopheles stephensi*. The antilog of X was then the LC₅₀ (conc. of 50% mortality) or LC₉₀ (conc. of 90% mortality) value.

The LC₅₀ & LC₉₀ were found to be 84.23ppm or 0.084v/v & 989.3956ppm or 0.989% v/v for *Anopheles stephensi*. One hundred percent mortality was observed for the concentrations tested against the organisms. The extract was effective against *Anopheles stephensi*.

Larvicidal activity of MECL against *Culex quinquefasciatus*

The results obtained for the larvicidal effect of MECL against *Culex quinquefasciatus* & the graphical representation were presented (Table 2 & Fig. 2).

From table 2, it can be observed that a mortality of 100.00 ± 0.00 was observed for *Culex quinquefasciatus*. The LC₅₀ and LC₉₀ values were calculated using Probit analysis. The percentage mortality calculated using Abbott's formula versus log concentration was plotted and Y=50 is substituted in the resulting linear equation to obtain the X value.

The linear regression equation was found to be $y = 44.533x - 26.161$ for activity against *Culex quinquefasciatus*. The antilog of X was then the LC₅₀ (conc. of 50% mortality) or LC₉₀ (conc. of 90% mortality) value. The LC₅₀ & LC₉₀ were found to be 51.31ppm or 0.051% v/v & 405.88ppm or 0.405% v/v for *Culex quinquefasciatus*. One hundred percent mortality was observed for the concentrations tested against the organisms. The extract was effective against *Culex quinquefasciatus*

Larvicidal activity of MECL against *Aedes aegypti*

The results obtained for the larvicidal effect of MECL against *Aedes aegypti* & the graphical representation were presented (Table 3 & Fig. 3).

From the Table 3, it was observed that a mortality of 100.00 ± 0.00 was observed for *Aedes aegypti*. The LC₅₀ and LC₉₀ values were calculated using Probit analysis. The percentage mortality calculated using Abbott's formula versus log concentration was plotted and Y=50 is substituted in the resulting linear equation to obtain the X value. The linear regression equation was found to be $y = 23.278x - 29.67$ for activity against *Aedes aegypti*. The antilog of X was then the LC₅₀ (conc. of 50% mortality) or LC₉₀ (conc. of 90% mortality) value.

The LC₅₀ & LC₉₀ were found to be 2645ppm or 2.645% v/v & 138326ppm or 138.326% v/v for *Aedes aegypti*. One hundred percent mortality was observed for the concentrations tested against the organisms. The extract was effective against *Aedes aegypti*. One hundred percent mortality was observed for various concentrations ranging for above organisms tested were observed within 24hr of the start of the experiment.

CONCLUSION

In the present study, Methanolic extract of *Citrullus lanatus* (MECL) showed promising larvicidal activity against important vectors of Malaria, Filariasis, Dengue and Yellow fever. MECL may be used as biodegradable, ecofriendly and cost effective alternative mosquito control tool instead of using synthetic chemicals.

Conflict of interest statement: We declare that we have no conflict of interest.

Table 1
Larvicidal activity of MECL against *Anopheles stephensi*

S. No	Conc. (ppm)	Log ₁₀ Conc.	Total No.	No. Dead	% Mortality	Probit	Probit	Regression value
0	0	0	0	0	0	0		0
1	25	1.40	20	3.00	15.00	3.77	3.00	4.358
2	50	1.70	20	5.00	25.00	4.19	4.00	4.778
3	100	2.00	20	6.00	30.00	4.36	4.00	4.948
4	150	2.18	20	9.00	45.00	4.8	5.00	5.388
5	200	2.30	20	10.00	50.00	4.92	5.00	5.508
6	250	2.40	20	13.00	65.00	5.33	6.00	5.918
7	300	2.48	20	16.00	80.00	5.81	6.00	6.398
8	350	2.54	20	17.00	85.00	5.99	6.00	6.578
9	400	2.60	20	19.00	95.00	6.65	7.00	7.238
10	450	2.65	20	20.00	100.00			
LC ₅₀			84.23ppm or 0.084v/v					
LC ₉₀			989.396ppm or 0.989% v/v					

Table 2
Larvicidal activity of MECL against *Culex quinquefasciatus*

S. No	Conc. (ppm)	Log ₁₀ Conc.	Total No.	No. Dead	% Mortality	Probit	Probit	Regression value
0	0	0	0	0	0	0	0	0.00
1	5	0.70	20	1.00	5.00	0	0.00	0.65
2	10	1.00	20	1.00	5.00	0	0.00	0.65
3	25	1.40	20	3.00	15.00	3.77	4.00	4.42
4	50	1.70	20	6.00	30.00	4.36	4.00	5.01
5	100	2.00	20	10.00	50.00	4.92	5.00	5.57
6	150	2.18	20	13.00	65.00	5.33	5.00	5.98
7	200	2.30	20	16.00	80.00	5.81	6.00	6.46
8	250	2.40	20	17.00	85.00	5.99	6.00	6.64
9	300	2.48	20	19.00	95.00	6.75	7.00	7.40
10	350	2.54	20	20.00	100.00			
LC ₅₀			51.31ppm or 0.051%v/v					
LC ₉₀			405.88ppm or 0.405%v/v					

Table 3
Larvicidal activity of MECL against *Aedes aegypti*

S. No	Conc. (ppm)	Log ₁₀ Conc.	Total No.	No. Dead	% Mortality	Probit	Probit	Regression value
0	0	0	0	0	0	0	0	0.00
1	5	0.70	20	1.00	5.00	0	0.00	0.65
2	10	1.00	20	1.00	5.00	0	0.00	0.65
3	25	1.40	20	3.00	15.00	3.77	4.00	4.42
4	50	1.70	20	6.00	30.00	4.36	4.00	5.01
5	100	2.00	20	10.00	50.00	4.92	5.00	5.57
6	150	2.18	20	13.00	65.00	5.33	5.00	5.98
7	200	2.30	20	16.00	80.00	5.81	6.00	6.46
8	250	2.40	20	17.00	85.00	5.99	6.00	6.64
9	300	2.48	20	19.00	95.00	6.75	7.00	7.40
10	350	2.54	20	20.00	100.00			
LC ₅₀			2645ppm or 2.645%v/v					
LC ₉₀			138326ppm or 138.326%v/v					




Fig. 1
Larvicidal activity of MECL against *Anopheles stephensi*




Fig. 2
Larvicidal activity of MECL against *Culex quinquefasciatus*




Fig. 3
Larvicidal activity of MECL against *Aedes aegypti*

REFERENCES

1. WHO. Fifteenth report of the WHO expert committee on vector biology and control- Vector resistance to pesticides, WHO Tech Rep Ser. 1992; 818: 1-62.
2. Arivoli S, Samuvel, Tennyson, Larvicidal efficacy of *Strychnus nuxvomica* Linn. (Family: Loganiaceae) leaf extracts against the synarial vector *Culex quinquefasciatus* (Diptera: Culicidae). World J of zoology, 2012; 1: 6-11.
3. Hadjiakhoondi A, Aghel N, Zamanizadeh N, Vatandoost H, Chemical and Biological study of *Mentha spicata* L essential oil from Iran. Daru, 2000; 8(1-2): 19-21.
4. Kumari R, Thapar BR, Dasgupta RK, Kaul SM, Lal S, Susceptibility status of Malaria vectors to insecticides in India. J Com Dis, 1998; 30(3): 179-185.
5. Davidson G, Alternative measures to insecticides for mosquito control. Pesticide Sci. 1972; 3: 503-514.
6. Deshmukh PB, Chavan SR, Renapurkar DM, A study of insecticidal activity of twenty indigenous plants. Pesticides, 1982; 16: 7-10.
7. Mittal P, Subbarao SK, Prospects of using herbal products in the control of mosquito vectors. ICMR Bull, 2003; 33(1): 1-10.
8. Oyolu CA, Quantitative and qualitative study of seed types in egusi (*Colocynthis citrullus* L). Tropic Sci, 1977; 19(1): 55-62.
9. Fursa TB, Intra specific classification of Watermelon under cultivation. Kulturpflanze 1981; 29(1): 297-300.
10. Aruna A, Vijayalakshmi K, Karthikeyan V, Nandhini SR, Jegadeesh S, Water Melon Plant (*Citrullus lanatus*): Pharmacognostical Standardization and Phytochemical Screening of its Leaves. Pharmanest, 2014; 5(4): 2184-2191.
11. Rahuman AA, Venkatesan P, Gopalakrishnan G, Mosquito larvicidal activity of oleic and linoleic acids isolated from *Citrullus colocynthis* (Linn.) Schrad. Parasitol Res, 2008; 103(6):1383-90.
12. Dharmagadda VSS, Naik SN, Mittal PK, Vasudevan P, Larvicidal activity of *Tagetes patula*. Bioresource Technol, 2005; 96: 1235-1240.
13. Singh RK, Mittal PK, Dhiman RC, Laboratory study on larvicidal properties of leaf extract of *Calotropis procera* (Family: Asclepiadaceae) against mosquito larvae. J Com Dis, 2005; 37: 109-113.
14. Thomas TG, Raghavendra K, Lal S, Saxena VK, Larvicidal properties of latex from unripe fruits of *Carica papaya* Linn (Caricaceae). J Com Dis, 2004; 36(4): 290-292.
15. Singh SP, Raghavendra K, Singh RK, Subbarao SK, Studies on the larvicidal properties of leaf extract of *Solanum nigrum* Linn (Family: Solanaceae). Curr Sci, 2001; 81(12): 1529-1530.
16. Mehra BK, Haridhar PK, The effect of crude acetone extract of *Annona squamosa* Linn (Family: Annonaceae) on possible control potential against larvae of *Culex quinquefasciatus* Say. J Entomol Res, 2001; 24(4): 141-146.
17. Thomas TG, Sharma SK, Jalees S, Rehman SJ, Larvicidal properties of an indigenous plant *Yucca aloifolia* Linn against mosquito larvae. J Basic Appl Bio Med, 1994; 2:53-55.
18. Pathak N, Mittal PK, Singh OP, Sagar V, Vasudevan P, Larvicidal action of essential oil from plants against the vector mosquitoes, *Anopheles stephensi* (Liston), *Culex quinquefasciatus* (Say) and *Aedes aegypti* (L). Int Pest Contr, 2000; 46: 53-55.
19. Hadjiakhoondi A, Sadeghipour-Roodsari HR, Vatandoost H, Khanavi M, Abai MR, Vosoughi M *et al*, Fatty acid composition and toxicity of *Melia azedarach* L. fruits against malaria vector *Anopheles stephensi*. Iran J Pharm Sci, 2006; 2(2): 97-102.